



AD735835

THE UNIVERSITY
OF WISCONSIN



MATHEMATICS RESEARCH CENTER




Address:

Mathematics Research
Center
The University of Wisconsin
Madison, Wisconsin 53706
U.S.A.

Reproduced by
NATIONAL TECHNICAL
INFORMATION SERVICE
Springfield, Va 22151

25

DISCLAIMER NOTICE



**THIS DOCUMENT IS BEST
QUALITY AVAILABLE. THE COPY
FURNISHED TO DTIC CONTAINED
A SIGNIFICANT NUMBER OF
PAGES WHICH DO NOT
REPRODUCE LEGIBLY.**

THE UNIVERSITY OF WISCONSIN
MATHEMATICS RESEARCH CENTER

Contract No. : DA-31-124-ARO-D-462


FIXED POINTS AND STABILITY FOR A SUM OF
TWO OPERATORS IN LOCALLY CONVEX SPACES

G. L. Cain, Jr. and M. Z. Nashed

This document has been approved for public
release and sale; its distribution is unlimited.

MRC Technical Summary Report #1115
September 1971

Received November 17, 1970



Madison, Wisconsin 53706

Security Classification

DOCUMENT CONTROL DATA - R & D		
(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)		
1. ORIGINATING ACTIVITY (Corporate author) Mathematics Research Center, University of Wisconsin, Madison, Wis. 53706		25. REPORT SECURITY CLASSIFICATION Unclassified
		26. GROUP None
3. REPORT TITLE FIXED POINTS AND STABILITY FOR A SUM OF TWO OPERATORS IN LOCALLY CONVEX SPACES		
4. DESCRIPTIVE NOTES (Type of report and inclusive dates) Summary Report: no specific reporting period.		
5. AUTHOR(S) (First name, middle initial, last name) G. L. Cain, Jr. and M. Z. Nashed		
6. REPORT DATE September 1971	7a. TOTAL NO. OF PAGES 21	7b. NO. OF REFS 28
8a. CONTRACT OR GRANT NO. Contract No. DA-31-124-ARO-D-462	8b. ORIGINATOR'S REPORT NUMBER(S) 1115	
8c. PROJECT NO. None	8d. OTHER REPORT NO(S) (Any other numbers that may be assigned this report) None	
10. DISTRIBUTION STATEMENT Distribution of this document is unlimited.		
11. SUPPLEMENTARY NOTES None	12. SPONSORING MILITARY ACTIVITY Army Research Office-Durham, N.C.	
13. ABSTRACT <p>Some fixed point theorems for a sum of two operators are proved, generalizing to locally convex spaces a fixed point theorem of M. A. Krasnoselskii, for a sum of a completely continuous and a contraction mapping, as well as some of its recent variants.</p> <p>A notion of stability of solutions of nonlinear operator equations in linear topological spaces is formulated in terms of specific topologies on the set of nonlinear operators, and a theorem on the stability of fixed points of a sum of two operators is given. As a byproduct, sufficient conditions for a mapping to be open or to be onto are obtained.</p>		

DD FORM 1473

Unclassified

Security Classification

AGO 8696A

ABSTRACT

Some fixed point theorems for a sum of two operators are proved, generalizing to locally convex spaces a fixed point theorem of M. A. Krasnoselskii, for a sum of a completely continuous and a contraction mapping, as well as some of its recent variants.

A notion of stability of solutions of nonlinear operator equations in linear topological spaces is formulated in terms of specific topologies on the set of nonlinear operators, and a theorem on the stability of fixed points of a sum of two operators is given. As a byproduct, sufficient conditions for a mapping to be open or to be onto are obtained.

FIXED POINTS AND STABILITY FOR A SUM OF TWO OPERATORS IN LOCALLY CONVEX SPACES

G. L. Cain, Jr. and M. Z. Nashed

1. Introduction

Several algebraic and topological settings in the theory and applications of nonlinear operator equations lead naturally to the investigation of fixed points of a sum of two nonlinear operators, or more generally, fixed points of a mapping on the Cartesian product $X \times X$ into X , where X is some appropriate space.

Fixed point theorems in topology and nonlinear functional analysis are usually based on certain properties (such as complete continuity, monotonicity, contractiveness, etc.) that the operator, considered as a single entity must satisfy. We recall for instance the Banach fixed point theorem, which asserts that a strict contraction on a complete metric space into itself has a unique fixed point, and the Schauder principle, which asserts that a continuous mapping F on a closed convex set K in a Hausdorff locally convex topological vector space X into K such that $F(K)$ is contained in a compact set, has a fixed point. In many problems of analysis, one encounters operators which may be split in the form $T = A + B$, where A is a contraction in some sense, and B is completely continuous, and T itself has neither of these properties. Thus neither the Schauder fixed point theorem nor the Banach fixed point theorem applies directly in this case, and it becomes

desirable to develop fixed point theorems for such situations. An early theorem of this type was given by Krasnoselskii [12]: Let X be a Banach space, S be a bounded closed convex subset of X , and A, B be operators on S into X such that $Ax + By \in S$ for every pair $x, y \in S$. If A is a strict contraction and B is continuous and compact, then the equation $Ax + Bx = x$ has a solution in S . The proof of this theorem is quite simple, given the Schauder principle.

Krasnoselskii's theorem is an example of an algebraic setting which leads to the consideration of fixed points of a sum of two operators. In this setting, a complicated operator is split into the sum of two simpler operators which have been well investigated and for which fixed point theorems abound. For recent contributions to fixed points of this type, see Remark 3.1.

There is another setting which also leads naturally to the investigation of fixed points of a sum of two operators. This setting arises from perturbation theory. Here the operator equation $Ax + Bx = x$ is considered as a perturbation of $Ax = x$ (or of $Bx = x$), and one would like to assert the existence of a solution of the perturbed equation, given that the original unperturbed equation has a solution. In such a setting, there is, in general, no continuous dependence of solutions on the perturbations. If, however, one requires such continuous dependence, then we have a general problem of stability of solutions, where stability is defined in terms of certain topologies on the class of operators under consideration.

The purpose of this paper is to prove some fixed point theorems in the two settings mentioned above.

2. Definitions and preliminaries

Throughout this paper, X will denote a Hausdorff locally convex topological vector space, and \mathcal{P} a family of seminorms which generates the topology of X . For $p \in \mathcal{P}$ and $r > 0$, the set $\{x \mid p(x - x_0) < r\}$ is denoted by $S_p(x_0, r)$. The closure of this set is denoted by $\bar{S}_p(x_0, r)$, and its boundary by $\partial S_p(x_0, r)$. We shall also sometimes use $V(p)$ to stand for $S_p(0, 1)$. A continuous mapping $F : X \rightarrow X$ is said to be p -completely continuous for $p \in \mathcal{P}$ if the closure of $F[\bar{S}_p(0, n)]$ is compact for each positive integer n .

Several generalizations of Schauder's fixed point theorem to locally convex topological vector spaces have been made by Tychonoff [26], Hukuhara [9], Yamamuro [28], Singhal [25], Nguyen-Xuan-Loc [17], and others. In the present paper, we shall be interested in the following variants of Schauder's fixed point theorem, which are listed in order of increasing generality.

Theorem 2.1. Let X be a Hausdorff locally convex topological vector space.

(a) Let K be a non-empty compact convex subset of X and let F be a continuous mapping of K into K . Then F has a fixed point in K .

(b) Let K be a non-empty closed convex set in X and let F be a continuous mapping of K into K such that $F(K)$ is contained in a compact set. Then F has a fixed point in K .

(c) Let F be a p -completely continuous mapping of X into X .

If F maps $\partial S_p(x_0, r)$ into $\bar{S}_p(x_0, r)$, then F has a fixed point in $\bar{S}_p(x_0, r)$.

Part (a) is simply Tychonoff's generalization of Schauder's theorem (for a proof, see Dunford and Schwartz [4] or Bonsall [1]). A simple and interesting proof of (b) is given by Singbal [25]. Part (c) is a generalization of Rothe's version of Schauder's theorem [22].

Let $D \subset X$ and $p \in \mathcal{P}$. A mapping $A : D \rightarrow D$ is said to be a p -contraction if there is a γ_p , $0 \leq \gamma_p < 1$, such that for all x, y in D , $p(Ax - Ay) \leq \gamma_p p(x - y)$.

Let \mathcal{u} be the neighborhood system of the origin obtained from \mathcal{P} . Then for any given $U \in \mathcal{u}$, there exist a finite number of seminorms in \mathcal{P} , say p_1, \dots, p_n , and $r_i > 0$, $i = 1, \dots, n$, such that $U = \bigcap_{i=1}^n r_i V(p_i)$, where $V(p) = \{x \mid p(x) < 1\}$.

Theorem 2.2. Suppose D is a sequentially complete subset of X and the mapping $A : D \rightarrow D$ is a p -contraction for every $p \in \mathcal{P}$. Then A has a unique fixed point \bar{x} in D , and $A^n x \rightarrow \bar{x}$ for every $x \in D$.

Proof. Let $x \in D$ and let $U = \bigcap_{i=1}^n r_i V(p_i)$ be given. For $y \in D$

and $k \geq 1$, we have

$$p_i(A^k y - y) \leq (1 - \gamma_i)^{-1} p_i(Ay - y).$$

Choose N so that for $m \geq N$,

$$\gamma_i^m (1 - \gamma_i)^{-1} p_i (Ax - x) \leq r_i, \quad i = 1, \dots, n.$$

Thus

$$\begin{aligned} p_i (A^{m+k} x - A^m x) &\leq (1 - \gamma_i)^{-1} p_i (A^{m+1} x - A^m x) \\ &\leq \gamma_i^m (1 - \gamma_i)^{-1} p_i (Ax - x) \leq r_i. \end{aligned}$$

Hence $\{A^k x\}$ is a Cauchy sequence in D and therefore converges to a point \bar{x} in D . Clearly $A\bar{x} = \bar{x}$, and the uniqueness of the fixed point follows as usual since X is Hausdorff.

3. Fixed points of a sum of two operators in locally convex spaces

We begin with a simple theorem which generalizes Krasnoselskii's fixed point theorem [12] to locally convex spaces.

Theorem 3.1. Let D be a convex and complete subset of X , and A, B be operators on D into X such that $Ax + By \in D$ for every pair $x, y \in D$. Suppose A is a p -contraction for every $p \in \mathcal{P}$, and B is continuous and $B(D)$ is contained in a compact set. Then there is a point \bar{x} in D such that $A\bar{x} + B\bar{x} = \bar{x}$.

Proof. For each $y \in D$, the mapping \tilde{A} defined by $\tilde{A}x = Ax + By$ is a p -contraction for each $p \in \mathcal{P}$ and maps D into D , so by Theorem 2.2, it has a fixed point, Ly . In other words, $Ly = \tilde{A}(Ly) = A(Ly) + By$. Thus for all u, v in D ,

$$Lu - Lv = A(Lu) - A(Lv) + Bu - Bv.$$

So for each $p \in P$, we have

$$p(Lu - Lv) \leq \gamma_p p(Lu - Lv) + p(Bu - Bv), \text{ or}$$

$$(3.1) \quad p(Lu - Lv) \leq (1 - \gamma_p)^{-1} p(Bu - Bv).$$

It is clear from (3.1) that the operator L is continuous. To see that $L(D)$ is contained in a compact set, let $\{Lx_a\}$ be a net in $L(D)$. Then $\{Bx_a\}$ has a convergent subnet $\{Bx'_a\}$, since $B(D)$ is contained in a compact set. Thus $\{Bx'_a\}$ is a Cauchy net, and by (3.1), so also is $\{Lx'_a\}$. Hence $L(D)$ is contained in a compact set, so L has a fixed point \bar{x} in D , and

$$\bar{x} = L\bar{x} = A(L\bar{x}) + B\bar{x} = A\bar{x} + B\bar{x}.$$

This completes the proof.

The various forms of the Schauder-Tychonoff theorem stated in Theorem 2.1 require a priori that a certain closed ball (or its boundary) be mapped into itself by the operator. In his work on integral equations, Dubrovskii [3] used an alternative approach of finding conditions on a completely continuous operator which guarantee the existence of some closed ball which is mapped into itself by the operator. In the next theorem, we use this technique in the setting of a sum of two operators to prove a fixed point theorem which contains as a special case a new variant of the Schauder-Tychonoff theorem in locally convex spaces. Before proceeding to the theorem, we shall give some needed definitions.

For an operator T , a point $x_0 \in X$, and a real number $r > 0$,
define for each $p \in \mathcal{P}$,

$$R_p(x_0, T, r) = r^{-1} \sup\{p(Tx - Tx_0) \mid p(x - x_0) \leq r\} \text{ and}$$

$$Q_p(x_0, T, a) = \{r \mid R_p(x_0, T, r) < a\}.$$

Now consider $Q_p(x_0, T, a)$ as a subset (possibly empty) of $[0, \infty]$, the one-point compactification of $[0, \infty)$, and let $\text{cl}(Q_p(x_0, T, a))$ denote the closure of $Q_p(x_0, T, a)$ relative to $[0, \infty]$. Define

$$\beta_p(x_0, T) = \inf\{a \mid \infty \in \text{cl}(Q_p(x_0, T, a))\}.$$

We shall say that T is p -quasibounded at x_0 if $\beta_p(x_0, T)$ exists. T is called quasibounded at x_0 if it is p -quasibounded at x_0 for each $p \in \mathcal{P}$.

Note that this notion of quasiboundedness generalizes that of Granas [8].

The following theorem generalizes Theorem 3 of Nashed and Wong [16].

Theorem 3.2. Suppose the mapping S is a p -contraction for every p in \mathcal{P} , with contraction constants γ_p , and suppose the mapping T is continuous and $\overline{T(X)}$ is compact. If X is complete and if there is an x_0 in X and a $p \in \mathcal{P}$ such that T is p -quasibounded at x_0 and

$$\gamma_p + \beta_p < 1,$$

then $(I - S - T)x = z$ always has a solution.

Proof. Choose a so that $\gamma_p + a < 1$ and $\infty \in \text{cl}(Q_p(x_0, T, a))$.
 Let $u_0 = (I - S - T)x_0$, and choose c so that $c > p(z - u_0)[1 - (\gamma_p + a)]^{-1}$
 and $c \in Q_p(x_0, T, a)$. Then $R_p(x_0, T, c) < a$. Now define the set

$$D = \{x \in X \mid p(x - x_0) \leq c\}.$$

It follows that for x and y in D , $Sx + Ty + z \in D$:

$$p(Sx + Ty + z - x_0) = p(Sx + Ty + z - u_0 - Sx_0 - Tx_0)$$

$$\leq p(Sx - Sx_0) + p(Ty - Tx_0) + p(z - u_0)$$

$$\leq \gamma_p c + ac + [1 - (\gamma_p + a)]c \leq c.$$

It now follows from Theorem 3.1 that there is an \bar{x} in D such that
 $S\bar{x} + T\bar{x} + z = \bar{x}$.

Remark 3.1. For various fixed point theorems for a sum of two operators in Banach and Hilbert spaces, see Krasnoselskii et al. [13], [14], Browder [2], Edmunds [5], Fučík [6], [7], Kirk [11], Nashed and Wong [16], Petryshyn [18], [19], Sadovskii [23], and Webb [27]. In some of this previous work, the theorems are formulated for a mapping $F(x, y)$, not necessarily of the form $Ax + By$. Nadler [15] considered mappings defined on the Cartesian product of two metric spaces which are contractions in one variable or in each variable separately and proved that under certain conditions, such mappings have fixed points.

Essentially the same proof as that of Theorem 3.1 yields the following result.

Theorem 3.1'. Let D be a convex and complete subset of X and suppose $F : D \times D \rightarrow D$ is such that for each $p \in P$, there is a constant γ_p , $0 \leq \gamma_p < 1$, so that

$$p(F(x, y) - F(x, z)) \leq \gamma_p p(y - z)$$

for all y, z in D . Suppose further that $B : D \rightarrow D$ is continuous, $B(D)$ is contained in a compact set, and

$$p(F(x, y) - F(z, y)) \leq p(Bx - Bz).$$

Then there is a point $\bar{x} \in D$ for which $F(\bar{x}, \bar{x}) = \bar{x}$.

Remark 3.2. Examining the proof of Theorem 3.1, one sees that if $D = \bar{S}_p(x_0, r)$, and X is complete, then we need only require that B be p -completely continuous. (We invoke 2.1c to obtain a fixed point of the operator L .)

For the operators considered in this section, the equation

$$(3.2) \quad Ax + Bx = x$$

has a solution in particular when A or B is the zero operator. Thus equation (3.2) may be considered as a perturbed equation associated with

$$(3.3) \quad Ax = x, \text{ or } Bx = x.$$

Theorems 3.1 and 3.2 state sufficient conditions under which the existence of a solution of either of the operator equations (3.3) is preserved under a perturbation of the operator in a certain class. We do not, however, have any information on how much of a change results in the solution. In particular, a slight perturbation of the operator A by an operator of type B need not necessarily produce only a slight change in the solution. In other words, in the algebraic setting of Theorems 3.1 and 3.2, one does not necessarily have continuous dependence of solutions of $Ax = x$ on perturbations of A by operators of the type B (or vice versa). We shall turn our attention in the next section to this question of continuous dependence of the solutions.

4. Stability of fixed points and solutions of nonlinear operator equations

In [10], Kasriel and Nashed formulated and investigated a notion of stability of solutions of some classes of nonlinear operator equations in Banach spaces in terms of specific topologies on the set of nonlinear operators, and obtained some results on the openness of certain mappings as a byproduct. In this section, we extend these formulations in several directions and prove a theorem on the stability of fixed points for the sum of two operators.

Let \mathcal{K} be a collection of continuous maps on X whose domains are such that if $A_0 \in \mathcal{K}$, $x_0 \in \text{domain of } A_0$, then $S_p(x_0, r) \subset \text{domain of } A_0$ for r sufficiently small. Let τ be a topology on \mathcal{K} . Suppose $A_0 \in \mathcal{K}$, $y_0 \in X$ and $A_0 x_0 = y_0$.

Definition 4.1. The solution x_0 of $A_0 u = y_0$ is called p -stable with respect to (K, \mathcal{T}) if for each $r > 0$ there exist $d > 0$ and a neighborhood Ω of A_0 such that for all $y \in S_p(y_0, d)$ and $A \in \Omega$, there exists an $x \in S_p(x_0, r)$ such that $Ax = y$. The solution x_0 is said to be a stable solution with respect to (K, \mathcal{T}) if it is p -stable solution for every $p \in P$.

For $A \in K$, (x_0, A, r) will be called a p -admissible triple if $\bar{S}_p(x_0, r)$ is contained in the domain of A .

Let K_p be the class of all continuous maps B from open subsets of X into X which are such that $I - B$ is p -completely continuous. If (x_0, B_0, r) is a p -admissible triple and $b > 0$, then $\Omega_U(x_0, B_0, r, p, b)$ will denote the collection of all $B \in K_p$ such that (x_0, B, r) is a p -admissible triple and $p(Bx - B_0 x) \leq b$ for all $x \in \bar{S}_p(x_0, r)$. Let \mathcal{T}_p be the topology on K_p generated by taking the collection of all such Ω_U as a subbase.

Now define

$$\tilde{R}_p(x_0, T, r) = r^{-1} \sup\{p(Tx - Tx_0) \mid p(x - x_0) = r\},$$

and

$$\eta_p(x_0, T) = \inf\{r \mid \tilde{R}_p(x_0, T, r) < 1\}.$$

Note that stability for the class K can be reduced to consideration of equations of the form $A_0 x = \theta$.

Theorem 4.1. Let $B_0 \in K_p$ and suppose $B_0 x_0 = \theta$. If $\eta_p(x_0, I - B_0) = 0$, then x_0 is a p -stable solution of $B_0 x = \theta$ with respect to (K_p, \mathcal{T}_p) .

Proof. Let $\epsilon > 0$ be given. There is an r , $0 < r < \epsilon$, such that $R = \tilde{R}_p(x_0, I - B_0, r) < 1$. Let a and d be positive numbers such that $a + d < (1 - R)r$. Let $B \in \Omega_U(x_0, B_0, r, p, a)$ and $y \in S_p(\theta, d)$. Consider the mapping F on $\bar{S}_p(x_0, r)$ defined by $Fx = x - Bx + y$.

Clearly F is p -completely continuous since $B \in \mathcal{K}_p$. If F maps $\partial S_p(x_0, r)$ into $\bar{S}_p(x_0, r)$, it has a fixed point $\bar{x} \in \bar{S}_p(x_0, r)$. Then $B\bar{x} = y$, with $\bar{x} \in \bar{S}_p(x_0, r) \subset S_p(x_0, \epsilon)$, which proves the theorem. Now we show that F indeed maps $\partial S_p(x_0, r)$ into $\bar{S}_p(x_0, r)$:

$$p(Fx - x_0) \leq p(x - B_0x - x_0) + p(Bx - B_0x) + p(y),$$

and

$$p(x - B_0x - x_0) \leq \tilde{R}_p(x_0, I - B_0, r)r = Rr.$$

Hence

$$p(Fx - x_0) \leq Rr + a + d \leq Rr + r - Rr = r.$$

If \mathcal{K}_C is the class of all continuous operators B from open subsets of X into X which are such that $I - B$ is completely continuous, and if \mathcal{T}_C is the topology on \mathcal{K}_C generated by taking as a subbase the sets $\Omega_U(x_0, B_0, r, p, b)$ for all $p \in \mathcal{P}$, then we have the following

Corollary. If $B_0 \in \mathcal{K}_C$ and $B_0x_0 = \theta$, and if $\eta_p(x_0, I - B_0) = 0$ for every $p \in \mathcal{P}$, then x_0 is a stable solution of $B_0x = \theta$ with respect to $(\mathcal{K}_C, \mathcal{T}_C)$.

We next turn our attention to the question of stability of sums of operators.

If $x_0 \in X$, A_0 is a continuous operator, and $U \in \mathcal{U}$, then we shall say (x_0, A_0, U) is an admissible triple if $x_0 + \overline{U} \subset \text{domain } A_0$. (Recall that \mathcal{U} is the neighborhood system of the origin obtained from \mathcal{P} .) Let \mathcal{C}_1 be the collection of all continuous operators A which are such that $I - A$ is a p -contraction for every $p \in \mathcal{P}$. (Hereafter called simply a contraction.) For A_0 in \mathcal{C}_1 , $p \in \mathcal{P}$, a and b real numbers, and (x_0, A_0, U) an admissible triple, we define $\Omega_1(x_0, A_0, U, p, a, b)$ to be the collection of all A in \mathcal{C}_1 such that

- i) (x_0, A, U) is an admissible triple,
- ii) $p((A - A_0)x - (A - A_0)x_0) \leq bp(x - x_0)$ for all $x \in x_0 + \overline{U}$,
- iii) $p(Ax_0 - A_0x_0) \leq a$.

We define π_1 to be the topology on \mathcal{C}_1 obtained by taking all such Ω_1 as a subbase.

Let \mathcal{C}_2 be the collection of all continuous operators B which are such that $I - B$ has its range contained in a compact set. For $B_0 \in \mathcal{C}_2$, $p \in \mathcal{P}$, r a real number, (x_0, B_0, U) an admissible triple, we define $\Omega_2(x_0, B_0, p, r)$ to be the collection of all $B \in \mathcal{C}_2$ such that

- i) (x_0, B, U) is an admissible triple, and
- ii) $p(Bx - Bx_0) \leq r$ for all $x \in x_0 + \overline{U}$.

We define \mathcal{T}_2 to be the topology on \mathcal{C}_2 with all such Ω_2 as a subbase.

Next let $C = C_1 \times C_2$ be the Cartesian product of C_1 and C_2 endowed with the product topology $\mathcal{T} = \mathcal{T}_1 \times \mathcal{T}_2$. Suppose K_0 is an operator such that $I - K_0 = S_0 + T_0$ for $(I - S_0, I - T_0) \in C$.

Definition 4.2. The solution x_0 of $K_0 u = y_0$ is called stable with respect to (C, \mathcal{T}) if for each $U \in \mathcal{U}$, there is a neighborhood Ω of $(I - S_0, I - T_0)$ and a $W \in \mathcal{U}$ such that for all $y \in y_0 + W$ and $(I - S, I - T) \in \Omega$, there exists an $x \in x_0 + U$ so that $Kx = y$, where $I - K = S + T$.

Recall the definition of $R_p(x_0, T_0, r)$ and $Q_p(x_0, T_0, a)$. For p in \mathcal{P} define

$$\alpha_p(x_0, T_0) = \inf\{a \mid 0 \in \text{cl}(Q_p(x_0, T_0, a))\}.$$

Theorem 4.2. Let X be complete. Suppose $K_0 x_0 = y_0$, where $I - K_0 = S_0 + T_0$ for $(I - S_0, I - T_0) \in C$. If $\gamma_p + \alpha_p < 1$ for every p in \mathcal{P} , then x_0 is a stable solution with respect to (C, \mathcal{T}) . (γ_p is p -contraction constant of S_0 and $\alpha_p \equiv \alpha_p(x_0, T_0)$.)

Proof. Once again we shall, without loss of generality, take $y_0 = \theta$. Let $U = \bigcap_{i=1}^n r_i V(p_i) \in \mathcal{U}$ be given. For each $i = 1, 2, \dots, n$, there is a $\xi_i > 0$ such that $\xi_i + \gamma_i < 1$ and $0 \in \text{cl}(Q_i(x_0, T_0, \xi_i))$, where γ_i denotes γ_{p_i} , etc. Choose $s_i \leq r_i$ so that $R_i(x_0, T_0, s_i) < \xi_i$. Now choose positive constants a_i, b_i, c_i, d_i , for each $i = 1, 2, \dots, n$, so that

$$b_i s_i + a_i + 2c_i + d_i < (1 - \xi_i - \gamma_i) s_i.$$

Let

$$B = I - T \in \bigcap_1^n \Omega_2(x_0, I - T_0, U, p_i, c_i),$$

and

$$A = I - S \in \bigcap_1^n \Omega_1(x_0, I - S_0, U, p_i, a_i, b_i).$$

Also, let $W = \bigcap_1^n d_i V(p_i)$.

Suppose $y \in W$ and consider $Sx + Tz + y$ for all x and z in $x_0 + U^*$, where $U^* = \text{cl}(\bigcap_1^n s_i V(p_i))$. We shall show that $Sx + Tz + y \in x_0 + U^*$:

$$\begin{aligned} Sx + Tz + y - x_0 &= Sx + Tz + y - S_0 x_0 - T_0 x_0 \\ &= (Sx - S_0 x_0) + (Tz - T_0 x_0) + y \\ &= (A - A_0)x - (A - A_0)x_0 + S_0 x - S_0 x_0 + (A - A_0)x_0 \\ &\quad + (Tz - T_0 z) + (T_0 x_0 - Tx_0) + (T_0 z - T_0 x_0) + y, \end{aligned}$$

where $A_0 = I - S_0$. Now for each $i = 1, 2, \dots, n$, we have

$$\begin{aligned} p_i(Sx + Tz + y - x_0) &\leq p_i((A - A_0)x - (A - A_0)x_0) + p_i(S_0 x - S_0 x_0) \\ &\quad + p_i((A - A_0)x_0) + p_i(Tz - T_0 z) + p_i(T_0 x_0 - Tx_0) + p_i(T_0 z - T_0 x_0) + p_i(y) \\ &\leq b_i p_i(x - x_0) + \gamma_i p_i(x - x_0) + a_i + c_i + c_i + R_i(x_0, T_0, s_i) s_i + p_i(y) \\ &\leq (1 - \xi_i - \gamma_i) s_i + (\gamma_i + \xi_i) s_i = s_i. \end{aligned}$$

So for every $x, z \in x_0 + U^*$, we have $Sx + Tz + y \in x_0 + U^*$; thus by Theorem 3.1, there is a point $\bar{x} \in x_0 + U^*$ so that $S\bar{x} + T\bar{x} + y = \bar{x}$, or $K\bar{x} = y$, where $I - K = S + T$.

Remark. If we take $T_0 = 0$ in Theorem 4.2, we get a stability theorem for the fixed point of a contraction mapping on a complete locally convex Hausdorff topological vector space X . We note, however, that it is possible to formulate other notions of "contraction" for which the fixed point is not necessarily stable. Let W_0 be an open neighborhood of $\theta \in X$, $x_0 \in X$, and $W = x_0 + W_0$. Let $F : W \rightarrow X$. We say that F is a weak contraction if there exists a convex, closed and bounded $V \subset W_0$ such that $x, y \in W$ and $y - x \in \lambda V$ imply $F(y) - F(x) \in \lambda \alpha V$ for some $0 < \alpha < 1$. Let F be a weak contraction on W into X , and $F(x_0) - x_0 \in (1 - \alpha)V$. Then there exists a unique fixed point \bar{x} of F , $\bar{x} \in x_0 + V$. However, this fixed point is obviously not necessarily stable.

5. Applications

The fixed point theorems of section 3 can be applied to obtain existence theorems for mixed nonlinear integral equations of Urysohn-Volterra and Hammerstein-Volterra types in locally convex spaces in the same manner as the fixed point theorem for a sum of two operators in Banach spaces were used in [16].

We now obtain as an application of Theorem 3.1, a sufficient condition for a mapping to be open, which generalizes conditions given in [10], [20], and [21]. Recall that a mapping $F : X \rightarrow Y$ is open at a point

$y_0 \in F(X)$ if y_0 is an interior point of $F(X)$; that is, if there is a neighborhood N of y_0 such that $N \subset F(X)$. It follows easily from Definition 4.2 that if $Ku = y_0$ has a stable solution with respect to (C, \mathcal{F}) , then K is open at y_0 . We can, however, find much weaker conditions which insure that K is open at y_0 . To this end, define

$$\varphi_p(x_0, T) = \inf\{a \mid Q_p(x_0, T, a) \neq \emptyset\},$$

and suppose K is as in section 4; that is, $I - K = S + T$ for $(I - S, I - T)$ in C .

Theorem 5.1. Assume X is complete. If $Kx_0 = y_0$ and for some p in \mathcal{P} it is true that $\gamma_p + \varphi_p < 1$, then K is open at y_0 .

Proof. We may without loss of generality take $y_0 = \theta$. Choose ξ so that $Q_p(x_0, T, \xi) \neq \emptyset$ and $\gamma_p + \xi < 1$. Let $s \in Q_p(x_0, T, \xi)$ and choose $d < (1 - \xi - \gamma_p)s$. We shall now show that $S_p(\theta, d)$ is contained in the range of K .

Let $u \in S_p(\theta, d)$ and consider $p(Sx + Ty + u - x_0)$ for x and y in $\bar{S}_p(x_0, s)$:

$$\begin{aligned} p(Sx + Ty + u - x_0) &= p(Sx + Ty + u - Sx_0 - Tx_0) \\ &\leq p(Sx - Sx_0) + p(Ty - Tx_0) + p(u) \\ &\leq \gamma_p s + \xi s + d < s. \end{aligned}$$

Thus by Theorem 3.1, there is an $\bar{x} \in \bar{S}_p(x_0, s)$ such that $S\bar{x} + T\bar{x} + u = \bar{x}$, which proves the theorem.

An immediate application of this result is the following theorem giving sufficient conditions for certain operators to be onto maps.

Theorem 5.2. Let $B : X \rightarrow X$ be a continuous operator such that $T(X)$ is contained in a compact set, where $T = I - B$. Suppose for each $x \in X$, there is a $p \in \mathcal{P}$ such that $\varphi_p(x, T) < 1$. Then the range of B is X .

Proof. B is open at each point of $B(X)$ from the previous theorem, so $B(X)$ is an open subset of X . We shall show that $B(X)$ is also a closed subset of X , and hence $B(X)$ must be all of the connected space X .

To show $B(X)$ is closed, let \bar{x} be an accumulation point of $B(X)$ and let $\{y_a\}$ be a net in $B(X)$ such that $y_a \rightarrow \bar{x}$. Let x_a be such that $Bx_a = y_a$. Then $\{Tx_a\}$ has a convergent subnet say $\{Tx'_a\}$. Since $Bx'_a = x'_a - Tx'_a$, and $\{Bx'_a\}$ and $\{Tx'_a\}$ converge, we then know that $\{x'_a\}$ converges. But $Bx'_a \rightarrow \bar{x}$, so $\bar{x} \in B(X)$. Thus $B(X)$ is closed, and the theorem is proved.

REFERENCES

1. F. E. Bonsall, Lectures on some fixed point theorems of functional analysis, Tata Institute of Fundamental Research, Bombay, India, 1962.
2. F. E. Browder, Fixed point theorems for nonlinear semicontractive mappings in Banach spaces, Arch. Rat. Mech. Anal. 21 (1966), 259-269.
3. W. Dubrovskii, Sur certaines équations intégrales nonlinéaires, U.S.S.R. Math. Zh. 30 (1939), 49-60.
4. N. Dunford and J. T. Schwartz, Linear operators, Vol. 1, Interscience Publishers Inc., New York, 1958.
5. D. E. Edmunds, Remarks on non-linear functional equations, Math. Ann. 174 (1967), 233-239.
6. S. Fučík, Fixed point theorems for a sum of nonlinear mappings, Comment. Math. Univ. Carolinae 9 (1968), 133-143.
7. ———, Solving of nonlinear operator equations in Banach space, Comment. Math. Univ. Carolinae 10 (1969), 177-186.
8. A. Granas, The theory of compact vector fields and some of its applications to topology of functional spaces I, Rozpr. Matematyczne 10 (1962), 1-93.
9. M. Hukuhara, Sur l'existence des points invariants d'une transformation dans l'espace fonctionnel, Japanese Journal Math. 20 (1950), 1-4.

10. R. H. Kasriel and M. Z. Nashed, Stability of solutions of some classes of nonlinear operator equations, Proc. Amer. Math. Soc. 17 (1966), 1036-1042.
11. W. A. Kirk, On nonlinear mappings of strongly semicontractive type, J. Math. Anal. Appl. 27 (1969), 409-412.
12. M. A. Krasnoselskii, Two remarks on the method of successive approximations, Uspehi Mat. Nauk 10 (1955), 123-127. (Russian)
13. _____, and P. P. Zabreiko, Construction of a new fixed point theorem, Dokl. Akad. Nauk SSSR 176 (1967) 1223. (Russian)
14. _____, _____, and R. I. Kachurovskii, On a fixed point theorem for operators in Hilbert space, Funktsional analiz i prilozhen 1, (1967), 93-94. (Russian)
15. S. Nadler, Sequences of contractions and fixed points, Pacific J. Math. 27 (1968), 579-585.
16. M. Z. Nashed, and J. S. W. Wong, Some variants of a fixed point theorem of Krasnoselskii and applications to nonlinear integral equations, J. Math. Mech. 18 (1969), 767-778.
17. Nguyen-Xuan-Loc, Fixed points and openness in a locally convex space, Proc. Amer. Math. Soc. 18 (1967), 987-991.
18. W. V. Petryshyn, Remarks on fixed point theorems and their extensions, Trans. Amer. Math. Soc. 126 (1967), 43-53.
19. _____, Fixed point theorems involving P-compact, semi-contractive, and accretive operators not defined on all of a Banach space, J. Math. Anal. Appl. 23 (1968), 335-354.

20. M. Reichback, Fixed points and openness, Proc. Amer. Math. Soc. 12 (1961), 734-736.
21. _____, Some theorems on mappings onto, Pacific J. Math. 10 (1960), 1397-1407.
22. E. H. Rothe, Zur Theorie der Topologischen Ordnung und der Vektorfelder in Banachschen Raumen, Compositio Math. 5 (1937), 177-197.
23. B. N. Sadovskii, On a fixed point theorem, Functional analiz i prilozhen 1 (1976), 74-76. (Russian)
24. J. Schauder, Der Fixpunktsatz in Funktionalraumen, Studia Math. 2 (1930) 171-180.
25. B. V. Singbal, Generalized form of Schauder-Tychonoff's fixed point principle, in F. E. Bonsall [1].
26. A. Tychonoff, Ein Fixpunktsatz, Math. Ann. 111 (1935), 767-776.
27. J. R. L. Webb, Fixed point theorems for nonlinear semicontractive operators in Banach spaces, J. London Math. Soc. (2) 1 (1969), 683-688.
28. S. Yamamuro, Some fixed point theorems in locally convex linear spaces, Proc. Japan Acad. 40 (1964), 1-12.

Mathematics Research Center

FIXED POINTS AND STABILITY FOR A
SUM OF TWO OPERATORS IN LOCALLY
CONVEX SPACES

G. L. Cain, Jr. and M. Z. Nashed

MRC Report No. 1115 AD

Contract No. : DA-31-124-ARO-D-462

September 1971
21 pp.

Some fixed point theorems for a sum of two operators are proved. A notion of stability of solutions is formulated in terms of specific topologies and a theorem on the stability of fixed points of a sum of two operators is given.

UNCLASSIFIED

Functional Analysis

Mathematics Research Center

FIXED POINTS AND STABILITY FOR A
SUM OF TWO OPERATORS IN LOCALLY
CONVEX SPACES

G. L. Cain, Jr. and M. Z. Nashed

MRC Report No. 1115 AD

Contract No. : DA-31-124-ARO-D-462

September 1971
21 pp.

Some fixed point theorems for a sum of two operators are proved. A notion of stability of solutions is formulated in terms of specific topologies and a theorem on the stability of fixed points of a sum of two operators is given.

UNCLASSIFIED

Functional Analysis

Mathematics Research Center

FIXED POINTS AND STABILITY FOR A
SUM OF TWO OPERATORS IN LOCALLY
CONVEX SPACES

G. L. Cain, Jr. and M. Z. Nashed

MRC Report No. 1115 AD

Contract No. : DA-31-124-ARO-D-462

September 1971
21 pp.

Some fixed point theorems for a sum of two operators are proved. A notion of stability of solutions is formulated in terms of specific topologies and a theorem on the stability of fixed points of a sum of two operators is given.

UNCLASSIFIED

Functional Analysis

Mathematics Research Center

FIXED POINTS AND STABILITY FOR A
SUM OF TWO OPERATORS IN LOCALLY
CONVEX SPACES

G. L. Cain, Jr. and M. Z. Nashed

MRC Report No. 1115 AD

Contract No. : DA-31-124-ARO-D-462

September 1971
21 pp.

Some fixed point theorems for a sum of two operators are proved. A notion of stability of solutions is formulated in terms of specific topologies and a theorem on the stability of fixed points of a sum of two operators is given.

UNCLASSIFIED

Functional Analysis

Mathematics Research Center

FIXED POINTS AND STABILITY FOR A
SUM OF TWO OPERATORS IN LOCALLY
CONVEX SPACES

G. L. Cain, Jr. and M. Z. Nashed

MRC Report No. 1115 AD

Contract No. : DA-31-124-ARO-D-462

Some fixed point theorems for a sum of two operators are proved. A notion of stability of solutions is formulated in terms of specific topologies and a theorem on the stability of fixed points of a sum of two operators is given.

UNCLASSIFIED

Functional Analysis

September 1971
21 pp.

Mathematics Research Center

FIXED POINTS AND STABILITY FOR A
SUM OF TWO OPERATORS IN LOCALLY
CONVEX SPACES

G. L. Cain, Jr. and M. Z. Nashed

MRC Report No. 1115 AD

Contract No. : DA-31-124-ARO-D-462

Some fixed point theorems for a sum of two operators are proved. A notion of stability of solutions is formulated in terms of specific topologies and a theorem on the stability of fixed points of a sum of two operators is given.

UNCLASSIFIED

Functional Analysis

September 1971
21 pp.

Mathematics Research Center

FIXED POINTS AND STABILITY FOR A
SUM OF TWO OPERATORS IN LOCALLY
CONVEX SPACES

G. L. Cain, Jr. and M. Z. Nashed

MRC Report No. 1115 AD

Contract No. : DA-31-124-ARO-D-462

Some fixed point theorems for a sum of two operators are proved. A notion of stability of solutions is formulated in terms of specific topologies and a theorem on the stability of fixed points of a sum of two operators is given.

UNCLASSIFIED

Functional Analysis

September 1971
21 pp.

Mathematics Research Center

FIXED POINTS AND STABILITY FOR A
SUM OF TWO OPERATORS IN LOCALLY
CONVEX SPACES

G. L. Cain, Jr. and M. Z. Nashed

MRC Report No. 1115 AD

Contract No. : DA-31-124-ARO-D-462

Some fixed point theorems for a sum of two operators are proved. A notion of stability of solutions is formulated in terms of specific topologies and a theorem on the stability of fixed points of a sum of two operators is given.

UNCLASSIFIED

Functional Analysis

September 1971
21 pp.